



























- Peng, Y. P., Chen, K. S., Lai, C. H., Lu, P. J., & Kao, J. H. (2006). Concentrations of H<sub>2</sub>O<sub>2</sub> and HNO<sub>3</sub> and O<sub>3</sub>-VOC-NO<sub>x</sub> sensitivity in ambient air in southern Taiwan. *Atmospheric Environment*, 40(35), 6741-6751.
- Pramono, G. H. (2008). Akurasi Metode IDW Dan Kriging Untuk Interpolasi Sebaran Sedimen Tersuspensi Di Maros, Sulawesi Selatan. *Forum Geografi*, 22(1), 145-158. (In Indonesian)
- Robinson, T.P., and Metternicht, G. (2006). Testing the performance of spatial interpolation techniques for mapping soil properties. *Computers and Electronics in Agriculture*, 50, 97-108
- Ryan, P. H., and Lemasters, G. K. (2007). NIH Public Access. *Inhalation Toxicology*, 19, 127-133.
- Tayanç, M. (2000). An assessment of spatial and temporal variation of sulfur dioxide levels over Istanbul, Turkey. *Environmental pollution*, 107(1), 61-9.
- Vaughan, G., Quinn, P. T., Green, a C., Bean, J., Roscoe, H. K., Van Roozendaal, M., & Goutail, F. (2006). SAOZ measurements of NO<sub>2</sub> at Aberystwyth. *Journal of environmental monitoring* 8(3), 353-61.
- Whitworth, K. W., Symanski, E., Lai, D., & Coker, A. L. (2011). Kriged and modeled ambient air levels of benzene in an urban environment: an exposure assessment study. *Environmental health* 10(1), 21